

LUK'YANOV, V.K.; FOKIN, A.G.

Geometrical effect in stripping reactions on deformed nuclei.
Izv. AN SSSR. Ser. fiz. 28 no.1:56-59 Ja '64. (MIRA 17:1)

1. Ob'yedinennyy institut yadernykh issledovaniy.

DZHAMALOV, O.B., doktor ekon. nauk; VOLOTKO, N.A.; YUN, D.N.,
kand. ekon. nauk; FOFONOV, B.M., kand. ekon. nauk;
KALIYAKIN, P.V., kand.ekon. nauk; DESYATCHIKOV, B.A.,
kand. ekon. nauk; KHUDKOVSKIY, A.B., kand. ekon. nauk;
ARTYKOV, A., kand. ekon. nauk; FOKIN, A.I.; UL'MASOV, A.,
kand. ekon. nauk; YAKOVENKO, Ye., red.; BAKHTIYAROV, A.,
tekhn. red.

[Principles of the economics of Uzbekistan industry] Osno-
vy ekonomiki promyshlennosti Uzbekistana; uchebnoe posobie
Tashkent, Gosizdat UzSSR, 1963. 282 p. (MIRA 17:1)

DESYATCHIKOV, B.A., kand. ekon. nauk; GABZAILOV, G.F., kand. ekon. nauk; KADYROV, Z., nauchn. sotr.; ABDUSHUKUROV, T.; KALYAKDI, P.V., kand. ekon. nauk; FOKIN, A.I., kand. ekon. nauk; BAKIYEVA, R.A., nauchn. sotr.; IBRAGIMOV, M., nauchn. sotr.; KARDASI, A.A., kand. ekon. nauk; KADANER, E.A.; NIKONOV, F.D., nauchn. sotr.; ANTONETS, G.M.; ARTYKOV, A.A., kand. ekon. nauk; TRUSOV, A.N.; OVCHAROVA, M.A., nauchn. sotr.; TSOY, P., nauchn. sotr.; KALYAKIN, P.V., kand. ekon. nauk, otv. red.; DZHAMALOV, O.B., doktor ekon. nauk, red.; ARTYKOV, A., kand. ekon. nauk, red.; DESYATCHIKOV, B.A., kand. ekon. nauk, red.; SHARIFKHODZHAYEV, M., kand. ekon. nauk, red.; DESYATNIK, F.M., red.; GOR'KOVAYA, Z.P., tekhn. red.

[Economics of the machinery manufacture of Uzbekistan] Ekonomika mashinostroeniia Uzbekistana. Tashkent, Izd-vo AN Uzb.SSR, 1963. 289 p. (MIRA 16:12)

1. Akademiya nauk Uzbekskoy SSR, Tashkent. Institut ekonomiki. (Uzbekistan--Machinery industry)

FOKIN, A.M.

Some character traits and scientific aspects of V.I.Vernadskii.
Och.p0 ist.geol.znan. no.11:7-20 '63. (MIRA 16:7)
(Vernadskii, Vladimir Ivanovich, 1863-1945)

FOKIN, A.M.; KUPARADZE, D.I.

Conditions governing the formation of secondary dispersion
halos and stray fluxes as revealed by the study of the
Dzamskoye iron ore deposit. Geol. sbor. [Kavk.] no.2:181-186
'62. (MIRA 17:1)

Translation from: Referativnyy zhurnal, Geologiya, 1957, Nr 1,
p 167 (USSR) 15-57-1-1048

AUTHORS: Fokin, A. M., Ramishvili, G. G.

TITLE: Soil Movements in Shida-Kakheti and the Role of Human
Factors (**Fazy razvitiya opolzney v Shida-Kakheti i
rol' antropogennogo faktora**)

PERIODICAL: Soobshch. AN GruzSSR, 1956, Vol 17, Nr 3, pp 213-217

ABSTRACT: In the region of Shida-Kakheti there occur strongly
fractured argillaceous conglomerates and sandy clays
of Miocene age, overlain by deluvial and loess-like
sandy loams. Half the precipitation falls from April
through June, strongly wetting the slopes. Three
phases of soil movements are distinguished. The first
belongs to the beginning of post-glacial time. The
base level of these movements was the first terrace
above the flood plain. These dislocations are in the

Card 1/3

15-57-1-1048

Soil Movements in Shida-Kakheti (Cont.)

nature of huge surficial movements. Traces of them have been preserved as elongated stabilized flowage depressions with areas up to two square meters. The second phase, ancient soil movements, belongs to the period between the eleventh and the sixteenth centuries. The base level for these movements was the flood plain. The age of the dislocations has been established by indirect evidence: the thickness of stumps, indications of skeletal trees, and the smoothness of terraces and mounds. Remains of ancient settlements are preserved in the zone of these movements. Forests and other vegetation have been cut down on the slopes and the soil has been cultivated. These practices have led to activation of the soil movements. The third phase of movements belongs to the present epoch. The movements have been rejuvenated only in the lower slopes because of stream erosion. Those districts utilized by man have also been affected. Individual foci of movement have arisen through moistening of the slopes because of inefficient drainage nets or of excavating for the narrow-gauge railway of the Akhmeta lumber

Card 2/3

15-57-1-1048

Soil Movements in Shida-Kakheti (Cont.)

establishment. The authors conclude that in the second and third phases the causes of the soil movements have been human activities. To fortify the slope, they recommend reforestation and strict regulation of irrigation systems.

Card 3/3

N. M. Ts.

MERABISHVILI, M.S., glavnyy red.; AVALIANI, G.A., red.; BAKRADZE, I.V., red.; DOLABERIDZE, L.D., red.; KAKABADZE, N.A., red.; KOMETIANI, G.A., red.; TVALCHRELIDZE, G.A., red.; TEGONIDZE, G.I., red.; FOKIN, A.M., red.; FILATOV, S.S., red.; EDILASHVILI, V.Ya., red.; BEREZOVSKAYA, L.I., red.izd-va; IVANOVA, A.G., tekhn.red.

[Yearbook of the Caucasus Institute of Raw Minerals for 1957]
Ezhegodnik Kavkazskogo instituta mineral'nogo syr'ia za 1957
god. Moskva, Gos.nauchno-tekhn.izd-vo lit-ry po geol. i okhrane
nedr, 1959. 54 p. (MIRA 13:12)

1. Tiflis. Kavkazskiy institut mineral'nogo syr'ya.
(Caucasus--Mines and mineral resources)

FOKIN, A.M.

Erosion belts on the southern slope of the Greater Caucasus
and their prospecting importance. Geol.sbor. [Kavk.] no.1:
125-131 '59. (MIRA 13:1)
(Caucasus--Erosion)

FOKIN, A.M.

Condenser for obtaining distilled water from industrial steam.
Sbor.rats.predl.vnedr.v proizvod. no.5:55 '60. (MIRA 14:8)

1. Moskovskiy trubnyy zavod.
(Condensers (Steam))

FOKIN, A.N.

Evolution of the supergene zone in the Chu-Ili Mountains.

Min.syr'e no.4:69-74 '62.

(MIRA 16:4)

(Chu-Ili Mountains—Weathering)

SOV/169-59-4-3476

Translation from: Referativnyy zhurnal, Geofizika, 1959, Nr 4, p 34 (USSR)

AUTHOR: Fokin, A.N.

TITLE:.. Using the IZh Device for Prospecting of Ore Deposits

PERIODICAL: Byul. nauchno-tekhn. inform. M-vo geol. i okhrany nedr SSSR, 1957, Nr 5 (10), pp 34 - 37

ABSTRACT: The IZh[✓] electric prospecting set may be used for discovering steeply dipping ore bodies of current-conducting and non-conducting character by the ratio-of-potentials method. This device is used successfully in areas of the difficulty classes I to V. It ensures a productivity of 1 km of electric profile per day when working with 5 m intervals. Having a weight of about 50 kg, the IZh set comprises a manually operated low frequency generator (80 - 100 cps) with an autotransformer producing a voltage of 50 - 400 v and a measuring bridge with an amplifier composed of 1K1P and 2P1P miniature tubes in the diagonal. The output is connected to headphones. The generator is connected to the AB feeder electrodes. A MON measuring set

Card 1/2

SOV/169-59-4-3476

Using the IZh Device for Prospecting of Ore Deposits

with three electrodes is used for determining the emf amplitude ratios between the central and each of the outer electrodes. The geological structure of the section in question is judged according to the value of this ratio. An efficient pulse generator composed of a phanotron has been developed to substitute the manually operated generator. Experimental models of this generator have been produced. Copper or steel tubes of 0.75 m length are used as electrodes. Plugs made of cotton cloth, impregnated by copper sulfate, are used on mountain slopes with widely evolved large block diluvium. The peculiarities of using the set under different prospecting conditions are described.

M.V. Sokol'skiy

Card 2/2

FOKIN, A.N.; SEMENOVA, G.A.; MILYAYEV, A.S.

Modern geological and geophysical methods of mapping weathering surfaces in prospecting for ore deposits as revealed by a study made in the arid-zone region. Kora vyvetr. no.6:272-282 '63.
(MIPA 17:9)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut mineral'nogo syr'ya, Moskva.

FOKIN, A.P.; PLANOVSKIY, A.N.; AKOPYAN, L.A.

Studying mass transfer during the drying of high-moisture content
products by means of atomizing in a uniflow apparatus. Plast. massy
no.8:43-44 '64. (MIRA 17:12)

FOKIN, A.P.; PLANOVSKIY, A.N.; AKOPYAN, L.A.

Calculation of spray dryers with allowance for stirring. Inzh.-
fiz. zhur. 8 no.1:116-118 Ja '65. (MIRA 18:3)

1. Institut khimicheskogo mashinostroyeniya, Moskva.

POKIN, A.S.

Giant-size lipoma of the breast. Sov.med.21 Supplement:16 '57.

(MIRA 11:2)

1. Iz khirurgicheskoy kliniki Saratovskogo meditsinskogo instituta.
(BREAST--TUMOR)

FOR 14, 11.5.

BUKOV, V.A., BYKOV, L.A., VALUK, V.A., VARTBARONOV, R.A., ZHILIS, E.F.,
KONDRAKOV, V.M., KUZ'MIN, V.A., SYCHEV, G.I. PROLOV, N.I.,
POKIN, A.S., KHARINSKIY, A.N. (Saratov)

New method for producing stable neurogenic hypertension in dogs
[with summary in English]. Arkh.pat. 20 no.5:21-27 '58 (MIRA 11:6)
(HEART, anatomy and histology,
thebesian vessels, review (Rus))

FORIN, A.S.

Effect of large doses of insulin on the content of sodium,
potassium, magnesium and total and ionized calcium in the
blood serum. Zhur. nevr. i psikh. 64 no.11:1717-1721 '64.
(MIRA 18:6)
1. Kafedra patologicheskoy fiziologii (zaveduyushchiy - prof.
N.I. Smutova) Leningradskogo pediatricheskogo meditsinskogo
instituta.

FOKINA, M.K.; FOKIN, A.S.

Experience in converting a library's holdings to the Universal
Decimal Classification. Opyt. rab. po tekhn. inform. i prop.
no.3:32-35 '63. (MIRA 16:12)

<p>1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50</p>																									
<p>A B C D E F G H I J K L M N O P Q R S T U V W X Y Z AA AB AC AD AE AF AG AH AI AJ AK AL AM AN AO AP AQ AR AS AT AU AV AW AX AY AZ BA BB BC BD BE BF BG BH BI BJ BK BL BM BN BO BP BQ BR BS BT BU BV BW BX BY BZ CA CB CC CD CE CF CG CH CI CJ CK CL CM CN CO CP CQ CR CS CT CU CV CW CX CY CZ DA DB DC DD DE DF DG DH DI DJ DK DL DM DN DO DP DQ DR DS DT DU DV DW DX DY DZ EA EB EC ED EE EF EG EH EI EJ EK EL EM EN EO EP EQ ER ES ET EU EV EW EX EY EZ FA FB FC FD FE FF FG FH FI FJ FK FL FM FN FO FP FQ FR FS FT FU FV FW FX FY FZ GA GB GC GD GE GF GG GH GI GJ GK GL GM GN GO GP GQ GR GS GT GU GV GW GX GY GZ HA HB HC HD HE HF HG HH HI HJ HK HL HM HN HO HP HQ HR HS HT HU HV HW HX HY HZ IA IB IC ID IE IF IG IH II IJ IK IL IM IN IO IP IQ IR IS IT IU IV IW IX IY IZ JA JB JC JD JE JF JG JH JI JJ JK JL JM JN JO JP JQ JR JS JT JU JV JW JX JY JZ KA KB KC KD KE KF KG KH KI KJ KL KM KN KO KP KQ KR KS KT KU KV KW KX KY KZ LA LB LC LD LE LF LG LH LI LJ LK LL LM LN LO LP LQ LR LS LT LU LV LW LX LY LZ MA MB MC MD ME MF MG MH MI MJ MK ML MN MO MP MQ MR MS MT MU MV MW MX MY MZ NA NB NC ND NE NF NG NH NI NJ NK NL NM NO NP NQ NR NS NT NU NV NW NX NY NZ OA OB OC OD OE OF OG OH OI OJ OK OL OM ON OO OP OQ OR OS OT OU OV OW OX OY OZ PA PB PC PD PE PF PG PH PI PJ PK PL PM PN PO PP PQ PR PS PT PU PV PW PX PY PZ QA QB QC QD QE QF QG QH QI QJ QK QL QM QN QO QQ QR QS QT QU QV QW QX QY QZ RA RB RC RD RE RF RG RH RI RJ RK RL RM RN RO RP RQ RR RS RT RU RV RW RX RY RZ SA SB SC SD SE SF SG SH SI SJ SK SL SM SN SO SP SQ SR SS ST SU SV SW SX SY SZ TA TB TC TD TE TF TG TH TI TJ TK TL TM TN TO TP TQ TR TS TT TU TV TW TX TY TZ UA UB UC UD UE UF UG UH UI UJ UK UL UM UN UO UP UQ UR US UT UU UV UW UX UY UZ VA VB VC VD VE VF VG VH VI VJ VK VL VM VN VO VP VQ VR VS VT VU VW VX VY VZ WA WB WC WD WE WF WG WH WI WJ WK WL WM WN WO WP WQ WR WS WT WU WV WW WX WY WZ XA XB XC XD XE XF XG XH XI XJ XK XL XM XN XO XP XQ XR XS XT XU XV XW XX XY XZ YA YB YC YD YE YF YG YH YI YJ YK YL YM YN YO YP YQ YR YS YT YU YV YW YX YY YZ ZA ZB ZC ZD ZE ZF ZG ZH ZI ZJ ZK ZL ZM ZN ZO ZP ZQ ZR ZS ZT ZU ZV ZW ZX ZY ZZ</p>																									
<p>1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50</p>																									
<p>A B C D E F G H I J K L M N O P Q R S T U V W X Y Z AA AB AC AD AE AF AG AH AI AJ AK AL AM AN AO AP AQ AR AS AT AU AV AW AX AY AZ BA BB BC BD BE BF BG BH BI BJ BK BL BM BN BO BP BQ BR BS BT BU BV BW BX BY BZ CA CB CC CD CE CF CG CH CI CJ CK CL CM CN CO CP CQ CR CS CT CU CV CW CX CY CZ DA DB DC DD DE DF DG DH DI DJ DK DL DM DN DO DP DQ DR DS DT DU DV DW DX DY DZ EA EB EC ED EE EF EG EH EI EJ EK EL EM EN EO EP EQ ER ES ET EU EV EW EX EY EZ FA FB FC FD FE FF FG FH FI FJ FK FL FM FN FO FP FQ FR FS FT FU FV FW FX FY FZ GA GB GC GD GE GF GG GH GI GJ GK GL GM GN GO GP GQ GR GS GT GU GV GW GX GY GZ HA HB HC HD HE HF HG HH HI HJ HK HL HM HN HO HP HQ HR HS HT HU HV HW HX HY HZ IA IB IC ID IE IF IG IH II IJ IK IL IM IN IO IP IQ IR IS IT IU IV IW IX IY IZ JA JB JC JD JE JF JG JH JI JJ JK JL JM JN JO JP JQ JR JS JT JU JV JW JX JY JZ KA KB KC KD KE KF KG KH KI KJ KL KM KN KO KP KQ KR KS KT KU KV KW KX KY KZ LA LB LC LD LE LF LG LH LI LJ LK LL LM LN LO LP LQ LR LS LT LU LV LW LX LY LZ MA MB MC MD ME MF MG MH MI MJ MK ML MN MO MP MQ MR MS MT MU MV MW MX MY MZ NA NB NC ND NE NF NG NH NI NJ NK NL NM NO NP NQ NR NS NT NU NV NW NX NY NZ OA OB OC OD OE OF OG OH OI OJ OK OL OM ON OO OP OQ OR OS OT OU OV OW OX OY OZ PA PB PC PD PE PF PG PH PI PJ PK PL PM PN PO PP PQ PR PS PT PU PV PW PX PY PZ QA QB QC QD QE QF QG QH QI QJ QK QL QM QN QO QQ QR QS QT QU QV QW QX QY QZ RA RB RC RD RE RF RG RH RI RJ RK RL RM RN RO RP RQ RR RS RT RU RV RW RX RY RZ SA SB SC SD SE SF SG SH SI SJ SK SL SM SN SO SP SQ SR SS ST SU SV SW SX SY SZ TA TB TC TD TE TF TG TH TI TJ TK TL TM TN TO TP TQ TR TS TT TU TV TW TX TY TZ UA UB UC UD UE UF UG UH UI UJ UK UL UM UN UO UP UQ UR US UT UU UV UW UX UY UZ VA VB VC VD VE VF VG VH VI VJ VK VL VM VN VO VP VQ VR VS VT VU VW VX VY VZ WA WB WC WD WE WF WG WH WI WJ WK WL WM WN WO WP WQ WR WS WT WU WV WW WX WY WZ XA XB XC XD XE XF XG XH XI XJ XK XL XM XN XO XP XQ XR XS XT XU XV XW XX XY XZ YA YB YC YD YE YF YG YH YI YJ YK YL YM YN YO YP YQ YR YS YT YU YV YW YX YY YZ ZA ZB ZC ZD ZE ZF ZG ZH ZI ZJ ZK ZL ZM ZN ZO ZP ZQ ZR ZS ZT ZU ZV ZW ZX ZY ZZ</p>																									

18

FOKIN, A.S.
ca

Processes and Properties Index

Hyposulfite. M. A. RABINOVICH and A. S. FOKIN. Russ. 23,379, Oct. 31, 1931.
The reduction of sulfites with Na-Hg obtained by electrolyzing NaCl is carried out in the presence of salts of silicic, boric or hydrofluoric acids as stabilizers.

ASM-SEA METALLURGICAL LITERATURE CLASSIFICATION

1930-1939

1940-1949

1950-1959

1960-1969

1970-1979

1980-1989

1990-1999

2000-2009

2010-2019

2020-2029

2030-2039

2040-2049

2050-2059

2060-2069

2070-2079

2080-2089

2090-2099

2100-2109

2110-2119

2120-2129

2130-2139

2140-2149

2150-2159

2160-2169

2170-2179

2180-2189

2190-2199

2200-2209

2210-2219

2220-2229

2230-2239

2240-2249

2250-2259

2260-2269

2270-2279

2280-2289

2290-2299

2300-2309

2310-2319

2320-2329

2330-2339

2340-2349

2350-2359

2360-2369

2370-2379

2380-2389

2390-2399

2400-2409

2410-2419

2420-2429

2430-2439

2440-2449

2450-2459

2460-2469

2470-2479

2480-2489

2490-2499

2500-2509

2510-2519

2520-2529

2530-2539

2540-2549

2550-2559

2560-2569

2570-2579

2580-2589

2590-2599

2600-2609

2610-2619

2620-2629

2630-2639

2640-2649

2650-2659

2660-2669

2670-2679

2680-2689

2690-2699

2700-2709

2710-2719

2720-2729

2730-2739

2740-2749

2750-2759

2760-2769

2770-2779

2780-2789

2790-2799

2800-2809

2810-2819

2820-2829

2830-2839

2840-2849

2850-2859

2860-2869

2870-2879

2880-2889

2890-2899

2900-2909

2910-2919

2920-2929

2930-2939

2940-2949

2950-2959

2960-2969

2970-2979

2980-2989

2990-2999

3000-3009

3010-3019

3020-3029

3030-3039

3040-3049

3050-3059

3060-3069

3070-3079

3080-3089

3090-3099

3100-3109

3110-3119

3120-3129

3130-3139

3140-3149

3150-3159

3160-3169

3170-3179

3180-3189

3190-3199

3200-3209

3210-3219

3220-3229

3230-3239

3240-3249

3250-3259

3260-3269

3270-3279

3280-3289

3290-3299

3300-3309

3310-3319

3320-3329

3330-3339

3340-3349

3350-3359

3360-3369

3370-3379

3380-3389

3390-3399

3400-3409

3410-3419

3420-3429

3430-3439

3440-3449

3450-3459

3460-3469

3470-3479

3480-3489

3490-3499

3500-3509

3510-3519

3520-3529

3530-3539

3540-3549

3550-3559

3560-3569

3570-3579

3580-3589

3590-3599

3600-3609

3610-3619

3620-3629

3630-3639

3640-3649

3650-3659

3660-3669

3670-3679

3680-3689

3690-3699

3700-3709

3710-3719

3720-3729

3730-3739

3740-3749

3750-3759

3760-3769

3770-3779

3780-3789

3790-3799

3800-3809

3810-3819

3820-3829

3830-3839

3840-3849

3850-3859

3860-3869

3870-3879

3880-3889

3890-3899

3900-3909

3910-3919

3920-3929

3930-3939

3940-3949

3950-3959

3960-3969

3970-3979

3980-3989

3990-3999

4000-4009

4010-4019

4020-4029

4030-4039

4040-4049

4050-4059

4060-4069

4070-4079

4080-4089

4090-4099

4100-4109

4110-4119

4120-4129

4130-4139

4140-4149

4150-4159

4160-4169

4170-4179

4180-4189

4190-4199

4200-4209

4210-4219

4220-4229

4230-4239

4240-4249

4250-4259

4260-4269

4270-4279

4280-4289

4290-4299

4300-4309

4310-4319

4320-4329

4330-4339

4340-4349

4350-4359

4360-4369

4370-4379

4380-4389

4390-4399

4400-4409

4410-4419

4420-4429

4430-4439

4440-4449

4450-4459

4460-4469

4470-4479

4480-4489

4490-4499

4500-4509

4510-4519

4520-4529

4530-4539

4540-4549

4550-4559

4560-4569

4570-4579

4580-4589

4590-4599

4600-4609

4610-4619

4620-4629

4630-4639

4640-4649

4650-4659

4660-4669

4670-4679

4680-4689

4690-4699

4700-4709

4710-4719

4720-4729

4730-4739

4740-4749

4750-4759

4760-4769

4770-4779

4780-4789

4790-4799

4800-4809

4810-4819

4820-4829

4830-4839

4840-4849

4850-4859

4860-4869

4870-4879

4880-4889

4890-4899

4900-4909

4910-4919

4920-4929

4930-4939

4940-4949

4950-4959

4960-4969

4970-4979

4980-4989

4990-4999

5000-5009

5010-5019

5020-5029

5030-5039

5040-5049

5050-5059

5060-5069

5070-5079

5080-5089

5090-5099

5100-5109

5110-5119

5120-5129

5130-5139

5140-5149

5150-5159

5160-5169

5170-5179

5180-5189

5190-5199

5200-5209

5210-5219

5220-5229

5230-5239

5240-5249

5250-5259

5260-5269

5270-5279

5280-5289

5290-5299

5300-5309

5310-5319

5320-5329

5330-5339

5340-5349

5350-5359

5360-5369

5370-5379

5380-5389

5390-5399

5400-5409

5410-5419

5420-5429

5430-5439

5440-5449

5450-5459

5460-5469

5470-5479

5480-5489

5490-5499

5500-5509

5510-5519

5520-5529

5530-5539

5540-5549

5550-5559

5560-5569

5570-5579

5580-5589

5590-5599

5600-5609

5610-5619

5620-5629

5630-5639

5640-5649

5650-5659

5660-5669

5670-5679

5680-5689

5690-5699

5700-5709

5710-5719

5720-5729

5730-5739

5740-5749

5750-5759

5760-5769

5770-5779

5780-5789

5790-5799

5800-5809

5810-5819

5820-5829

5830-5839

5840-5849

5850-5859

5860-5869

5870-5879

5880-5889

5890-5899

5900-5909

5910-5919

5920-5929

5930-5939

5940-5949

5950-5959

5960-5969

5970-5979

5980-5989

5990-5999

6000-6009

6010-6019

6020-6029

6030-6039

6040-6049

6050-6059

6060-6069

6070-6079

6080-6089

6090-6099

6100-6109

6110-6119

6120-6129

6130-6139

6140-6149

6150-6159

6160-6169

6170-6179

6180-6189

6190-6199

6200-6209

6210-6219

6220-6229

6230-6239

6240-6249

6250-6259

6260-6269

6270-6279

6280-6289

6290-6299

6300-6309

6310-6319

6320-6329

6330-6339

6340-6349

6350-6359

6360-6369

6370-6379

6380-6389

6390-6399

6400-6409

6410-6419

6420-6429

6430-6439

6440-6449

6450-6459

6460-6469

6470-6479

6480-6489

6490-6499

6500-6509

6510-6519

6520-6529

6530-6539

6540-6549

6550-6559

6560-6569

6570-6579

6580-6589

6590-6599

6600-6609

6610-6619

6620-6629

6630-6639

6640-6649

6650-6659

6660-6669

6670-6679

6680-6689

6690-6699

6700-6709

6710-6719

6720-6729

6730-6739

6740-6749

6750-6759

6760-6769

6770-6779

6780-6789

6790-6799

6800-6809

6810-6819

6820-6829

6830-6839

6840-6849

6850-6859

6860-6869

6870-6879

6880-6889

6890-6899

6900-6909

6910-6919

6920-6929

6930-6939

6940-6949

6950-6959

6960-6969

6970-6979

6980-6989

6990-6999

7000-7009

7010-7019

7020-7029

7030-7039

7040-7049

7050-7059

7060-7069

7070-7079

7080-7089

7090-7099

7100-7109

7110-7119

7120-7129

7130-7139

7140-7149

7150-7159

7160-7169

7170-7179

7180-7189

7190-7199

7200-7209

7210-7219

7220-7229

7230-7239

7240-7249

7250-7259

7260-7269

7270-7279

7280-7289

7290-7299

7300-7309

7310-7319

7320-7329

7330-7339

7340-7349

7350-7359

7360-7369

7370-7379

7380-7389

7390-7399

7400-7409

7410-7419

7420-7429

7430-7439

7440-7449

7450-7459

7460-7469

7470-7479

7480-7489

7490-7499

7500-7509

7510-7519

7520-7529

7530-7539

7540-7549

7550-7559

7560-7569

7570-7579

7580-7589

7590-7599

7600-7609

7610-7619

7620-7629

7630-7639

7640-7649

7650-7659

7660-7669

7670-7679

7680-7689

7690-7699

7700-7709

7710-7719

7720-7729

7730-7739

7740-7749

7750-7759

7760-7769

7770-7779

7780-7789

7790-7799

7800-7809

7810-7819

7820-7829

7830-7839

7840-7849

7850-7859

7860-7869

7870-7879

7880-7889

7890-7899

7900-7909

7910-7919

7920-7929

7930-7939

7940-7949

7950-7959

7960-7969

7970-7979

7980-7989

7990-7999

8000-8009

8010-8019

8020-8029

8030-8039

8040-8049

8050-8059

8060-8069

8070-8079

8080-8089

8090-8099

8100-8109

8110-8119

8120-8129

8130-8139

8140-8149

8150-8159

8160-8169

8170-8179

8180-8189

8190-8199

8200-8209

8210-8219

8220-8229

8230-8239

8240-8249

8250-8259

8260-8269

8270-8279

8280-8289

8290-8299

8300-8309

8310-8319

8320-8329

8330-8339

8340-8349

8350-8359

8360-8369

8370-8379

8380-8389

8390-8399

8400-8409

8410-8419

8420-8429

8430-8439

8440-8449

8450-8459

8460-8469

8470-8479

8480-8489

8490-8499

8500-8509

8510-8519

8520-8529

8530-8539

8540-8549

8550-8559

8560-8569

8570-8579

8580-8589

8590-8599

8600-8609

8610-8619

8620-8629

8630-86

1ST AND 2ND ORDER		PROCESS AND PROPERTIES INDEX		3RD AND 4TH ORDER	
<p>FOKIN, A. S.</p> <p>BC</p> <p>DIY</p> <p>Electrochemical production of sodium hydroxide. U. M. A. Ramanovitch and A. S. Fokin. B. P. E. Suvorovskii, A. P. Manovarov, and A. S. Fokin. <i>IZV. P. B. Suvorovskii (Ussr. Chem. J. 1931, 4, (Tech.), 204-205, 206-211, 212-219).</i>—1. SO₂ is reduced at a Hg-Na cathode, in which Na is removed electrolytically. The yield of NaOH is 70-80% of theory, and any desired concentration can be obtained. The chief source of loss is due to the reaction: $2\text{Na}_2\text{O}_2 \rightarrow \text{Na}_2\text{O}_2 + \text{Na}_2\text{O}_2$. Aq. Na₂O₂ is stabilized by Na₂SO₄.</p>					
<p>ASS-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>					
FROM STUDYING		FROM BOWING		FROM BOWING	
100000 01		100000 01		100000 01	
100000 01		100000 01		100000 01	

Fokin, A. S.

USSR/Chemical Technology. Chemical Products and Their Application--Synthetic fibers. I-26

Abs Jour: Ref Zhur-Khimiye, No 3, 1957, 10083

Author : Fokin, A. S.

Inst : Kiev Technical Institute of the Light Industry

Title : The Viscosity of Concentrated Gelatin Solutions at Elevated Temperatures

Orig Pub: Tr. Kievsk. tekhnol. in-ta legkoy prom-sti, 1955, No 7, 33-43

Abstract: Gelatin solutions containing 15% protein at 40° have a structural viscosity determined by the space lattice formed by the anisodiametric protein molecules. The energy of formation of that lattice apparently is not very large. Samples of commercial gelatine and proteins extracted from leather wastes have been investigated. It has been established that solutions of proteins extracted by neutralization from chrome-tanning

Card 1/2

Problem of the measurement of a period in the
of metals by means of a ...

91

FOKIN, A. V.

FOKIN, A.V.: "Investigation of the movement of the comet Oterma 3,
July 1942." Acad Sci USSR. Main Astronomical Observatory. Leningrad,
1956. (DISSERTATION For the Degree of Candidate in PHYSICOMATHEMATICAL
SCIENCE.)

So: Knizhaya letopis', No. 24, 1956

FOKIN, A.V.

Orbit of Oterma's comet 3 (1942 VII) according to observations
in 1943-1950. Biul. Inst. teor. astron. 7 no.2:89-112 '58.

(MIRA 13:3)

(Comet, Oterma's)

POKIN, A.V.

Investigating the close approach of Oterma's comet 3 to Jupiter.
Biul. Inst. teor. astron. 7 no.2:113-119 '58. (MIRA 13:3)
(Comet, Oterma's) (Jupiter (Planet))

3(1)

AUTHOR: Fokin, A.V.

SOV/33-35-4-23/25

TITLE: An Investigation of the Movement of the Comet Oterma 3 1942 VII (Issledovaniye dvizheniya komety oterma 3 1942 VII)

PERIODICAL: Astronomicheskii zhurnal, 1958, Vol 35, Nr 4, pp 674-676(USSR)

ABSTRACT: The comet Oterma 3 discovered on April 3, 1943 approached Jupiter a short time before the discovery (1937-1939). The author uses 127 observations of the years 1943-1950 in order to set up the differential equations of the baricentric motion of the comet, in order to integrate them then with intervals of 20 days back to December 1939. Furthermore the author applied planetocentral coordinates. Thus it was stated that the comet moved in the radius of action of Jupiter from December 27, 1936 to December 15, 1938 and that before this time the orbit of the comet was totally between the orbits of Jupiter and of Saturn, and therefore it could not be observed. Near the Jupiter the orbit was completely changed. The next Jupiter contiguity will be in 1961.

Card 1/2

An Investigation of the Movement of the Comet
Oterma 3 1942 VII

SOV/33-35-4-23/25

There are 2 tables, and 3 references, 1 of which is Soviet,
and 2 American.

ASSOCIATION: Odesskoye vyssheye morekhodnoye uchilishche (Odessa Higher
Navigation School)

SUBMITTED: March 14, 1957

Card 2/2

S/035/62/000/011/008/079
A001/A101

AUTHOR: Fokin, A. V.

TITLE: Ascending motion of a rocket in the gravity field

PERIODICAL: Referativnyy zhurnal, Astronomiya i Geodeziya, no. 11, 1962, 10,
abstract 11A77 ("Byull. In-ta teor. astron. AN SSSR", 1962, v. 8,
no. 5, 335 - 342, English summary)

TEXT: The author considers the problem of the motion of a rocket, whose mass varies according to an exponential law, in the Earth's gravity field. The latter is assumed to be central and the rocket motion proceeding along the radius-vector. Air resistance is neglected. The equation of the rocket motion is integrated, and the solution is represented in terms of elliptic integrals of the first and second kind. The sequence of calculations by the formulae obtained is presented.

Yu. B.

[Abstracter's note: Complete translation]

Card 1/1

ACCESSION NR: AT4001203

S/2511/62/008/005/0335/0342

AUTHOR: Fokin, A. V.

TITLE: Ascending motion of rocket in a gravitational force field

SOURCE: AN SSSR. Inst. teor. astron. Byulleten', v. 8, no. 5, 1962, 335-342

TOPIC TAGS: material point, earth gravitational force, reactive acceleration, gravitational acceleration, rocket, Kabal'chiah number, variable mass point, rocket motion, earth gravitational field, gravitational force field, rocket ascent, elliptic integral

ABSTRACT: The ascending motion in the earth's gravitational field of a rocket whose mass varies exponentially is calculated without allowance for the resistance of the medium. This problem was solved originally by A. A. Kosodem'yanskiy (Uch. zap. MGU, 154, Mekhanika, 4) in terms of elementary functions. The present solution is in

Card 1/2

ACCESSION NR: AT4001203

terms of elliptic integrals of the first and second kind, and the availability of very accurate tables for these integrals makes the use of the elliptic solution as convenient as calculations with elementary functions. Orig. art. has: 52 formulas.

ASSOCIATION: Inst. Teor. Astron. AN SSSR (Institute of Theoretical Astronomy AN SSSR)

SUBMITTED: 22Apr61

DATE ACQ: 21Nov63

ENCL: 00

SUB CODE: GM, AS

NO REF SOV: 001

OTHER: 000

Card 2/2

PHASE I BOOK EXPLOITATION

SOV/6488

Knunyants, Ivan Lyudvigovich, Academician, and Aleksandr Vasil'yevich Fokin, Professor

Pokoreniye nepristupnogo elementa (Conquest of an Inaccessible Element) Moscow, Izd-vo AN SSSR, 1963. 189 p. (Series: Akademiya nauk SSSR. Nauchno-populyarnaya seriya) Errata printed on inside of back cover. 25,000 copies printed.

Ed. of Publishing House: V. M. Tarasenko; Tech. Ed.: S. P. Golub'.

PURPOSE: This textbook is intended for chemists and engineers.

COVERAGE: The book covers the full range of fluorine chemistry and technology and is based on Soviet and Western sources. The text includes: properties of fluorocarbons; fluorinated hydrocarbons, e.g., fluorinated olefins which have high thermal stability and chemical resistance; esters of dibasic

Card 1/5
3

Conquest of an Inaccessible Element

SOV/6488

perfluorocarboxylic acids which are used in the preparation of stable lubricants and in synthetic rubber; preparation, properties, and applications of fluorine-containing plastics [teflon, teflon-100, poly(vinylidene fluoride)]; their radiation polymerization; fluorine containing elastomers used in supersonic aircraft and jet engines; freons; and other applications. The fifth chapter is devoted to lubricants and hydraulic fluids with high thermal and oxidation stability, which can be used under severe conditions and in systems with aggressive media. It is noted that chlorofluorocarbon oils lower the friction coefficient of rubbing surfaces made of the same metal under heavy loads and at high temperatures. Certain Soviet oils, greases, and hydraulic fluids are also described. Tabulated data are given for: 4-F, 3-F, 3-OK, 10-OK, 20-F, Summer No.5, Winter No.8, UPI, Fluid No.12, manometric fluid, and balancing fluid. Only KS, UPI, and Summer No.5 are insoluble in ammonia and amines. In describing esters of dicarboxylic acids and fluorinated

Card 2/5

Conquest of an Inaccessible Element

SOV/6488

alcohols, the author mentions the applicability of such lubricants for submarines to avoid a detectable oil film on the water surface (the ONR research is quoted). Modern lubricants are already in use in jet engines and other engines working at high temperatures. The author points out the importance of fluorine compounds in rocketry, aviation, astronautics, atomic energy, and industry. The last chapter is devoted to fluorine organic compounds used for: the improvement of light fastness of dyes (benzotri-fluoride, fluorobenzene, trifluoromethylbenzene); the preparation of chemotherapeutic compounds; and the preparation of a class of toxic compounds (fluoroacetates). These compounds belong to esters of the general type $F(Ch_2)_n COOR$, fluoroalcohols, and fluoroacetamides with their derivatives. Fluorophosphates are mentioned as CW compounds (e.g., DFP-3, Sarin, Soman) which caused a change in anti-CV methods. No personalities are mentioned. There are no references.

Card 3/3
/3

KNUNYANTS, I.L.; FOKIN, A.V.; KOSYREV, Yu.M.; SOROCHKIN, I.N.; FROSINA, K.V.

Nitration of perfluorobutadiene with nitrogen peroxide. Izv. AN SSSR
Ser.khim. no. 10:1772-1775 0 '63. (MIRA 17:3)

FOKIN, A.V., kand. fiziko-matematicheskikh nauk dotsent; KONONENKO, V.V., inzh.

Simple method for the determination of mean indicated pressure
by developed indicator diagrams. Izv. vys. ucheb. zav.;
mashinostr. no.2:71-75 '64. (MIRA 17:5)

1. Odesskoye vyssheye inzhenerno-morakoye uchilishche.

FOKIN, A.V.; KRINETSKIY, I.I.

Some general problems of the invariance theory. Avtom. upr. i vych. tekhn.
no.6:175-182 '64. (MIRA 17:10)

L 21735-65 DWT(m)/EPF(c)/EPR/EXP(j) Pc-Li/Pr-Li/Ps-Li SSD(a)/RPL RM/WW

ACCESSION NR: AP4044703

S/0062/64/000/008/1425/1429 13

AUTHOR: Knunyants, I. L.; Dyatkin, B. L.; Fokin, A. V.; Komarov, V. A.

TITLE: Nitration of perfluoroisobutylene¹

SOURCE: AN SSSR. Izvestiya. Seriya khimicheskaya, no. 8, 1964, 1425-1429

TOPIC TAGS: perfluoroisobutylene, nitration, nitrogen tetroxide reaction, nitroperfluorobutylnitrite, perfluoroisobutyldinitrite, nitroperfluorobutanol, bistrifluoromethylglycolic acid

ABSTRACT: Perfluoroisobutylene was heated with an equimolecular amount of N_2O_4 in a steel autoclave at 170-180°C for 6-8 hours to attain nearly complete conversion. Nitroperfluoro-*tert.*-butyl nitrite (compound VI in the literature), boiling 100-110°C, and a fraction boiling 30-100°C, apparently a mixture of perfluoroisobutyl- α, β -dinitrite, $(\text{CF}_3)_2\text{C}(\text{ONO})\text{CF}_2\text{ONO}$, and its conversion products were obtained. Hydrolysis of the 30-100°C fraction gave bistrifluoromethylglycolic acid (X) in 27% yield based on initial perfluoroisobutylene. Nitroperfluoro-

Cor 3/3

L 21735-65

ACCESSION NR: AP4044703

tert-butanol (IX) was obtained in 23% yield, based on initial perfluoroisobutylene, /
analysis of VI. No dinitro compound $(CF_3)_2C(NO_2)CFNO_2$ nor nitronitrite
 $(CF_3)_2C(NO_2)CF_2ONO$, nor products which could be obtained by their conversion
was found, contrary to earlier data by I. L. Knunyants and A. V. Fakin
(Dokl. Akad. Nauk SSSR 111, 1035 (1956)). Orig. art. has: 7 equations and 10 formulae

ASSOCIATION: Institut elementoorganicheskikh soyedineniy Akademii nauk SSSR
(Institute of Organometallic Compounds, Academy of Sciences, SSSR)

SUBMITTED: 28Dec62

ENCL: 01

SUB CODE: GC, MT

NO REF SOV: 005


OTHER: 004

Card 2/3

L 21735-65

ACCESSION NR: AP4044703

ENCLOSURE: 01



Card 3/3

FOKIN, A.V.; KOMAROV, V.A.; SKIADNEV, A.A.; DAVYDOVA, S.M.

Reactivity of nitroperfluoroalkyl nitrites and products of their transformation. Part 1: Reaction of nitroperfluoroalkyl nitrites with hydrogen sulfide. Zhur. ob. khim. 35 no.9:1662-1664 8 166

Reactivity of nitroperfluoroalkyl nitrites and products of their transformation. Part 2: Reaction of nitroperfluoroalkyl nitrites with mercaptans. Ibid.:1664-1666 (MIRA 18:10)

L 32682-66 EWT(m)/EWP(j) RM/FDN/JW

ACC NR: AP6012527

SOURCE CODE: UR/0062/66/000/003/0466/0472

AUTHOR: Knunyants, I. L.; Fokin, A. V.; Komarov, V. A.

ORG: none

TITLE: Nitration¹ of perfluoropropylene¹ with nitrogen dioxide and investigation of nitration products

SOURCE: AN SSSR. Izvestiya. Seriya khimicheskaya, no. 3, 1966, 466-472

TOPIC TAGS: nitration, organic chemistry, nitrogen oxide, fluorine compound, PROPYLENE

ABSTRACT: The present study is a continuation of work reported in *Dokl. AN SSSR*, III, 1035 (1956). The synthesized nitration products are given in the following table along with some of their properties:

UDC: 542.958.1 + 661.723-16

Card 1/3

L 32682-66

ACC NR: AP6012527

Formula	Boiling point °C (pres- sure, mm Hg)	d_4^{20}	n_D^{20}	Formula	Boiling point °C (pres- sure, mm Hg)	d_4^{20}	n_D^{20}
$\text{CF}_3\text{--CF}_2\text{--CF}_2\text{--ONOONO}_2$	57	1,637	1,3276	$\text{CF}_3\text{--C--CF}_2\text{NO}_2$ $\text{OH} \quad \text{OC}_2\text{H}_5$	118,5	1,391	1,3520
$\text{CF}_3\text{--C--CF}_2\text{NO}_2$ $\text{OH} \quad \text{OH}$	119-120	1,638	1,3560	CF_3NO_2 C--CH_3 O--C=O	68(44)	1,616	1,3621
$\text{CF}_3\text{--C--CF}_2\text{NO}_2$ O	32-33	1,5350	1,2955	CHF_2NO_2	42-43	1,4605	1,3158
$\text{CF}_3\text{--C--CF}_2\text{NO}_2$ $\text{OH} \quad \text{Cl}$	37	1,609	1,3500	$\text{CF}_3\text{--CH--CH}_3$ $\text{NO}_2 \quad \text{OH}$	64-65(25)	1,390	1,3825
$\text{CF}_3\text{--C--CF}_2\text{NO}_2$ $\text{OH} \quad \text{Br}$	50	1,935	1,3758	$\text{CF}_3\text{--CH}_2\text{--OH}$	65(40)	1,4792	1,3780
$\text{CF}_3\text{--C--CF}_2\text{NO}_2$ $\text{OH} \quad \text{CN}$	64-65(20)	1,6282	1,3405	NO_2 CH_3 $\text{CF}_3\text{--C--CH}_3$ $\text{NO}_2 \quad \text{OH}$	55(35)	1,2950	1,3015


During nitration of perfluoropropylene with nitrogen dioxide, nitroperfluoroisopropyl nitrile and dinitroperfluoropropane form. Hydrolysis of nitroperfluoroisopropyl nitrite produces nitroperfluoroacetone hydrate which upon dehydration produces anhydrous

Cord 2/3

L 32682-66

ACC NR: AP6012527

nitroperfluoroacetone. The chemical properties of nitroperfluoroacetone were investigated and some of its derivatives were synthesized. Nitroperfluoroacetone reacts with basic compounds to form difluoronitromethane and trifluoroacetic acid derivatives. The reaction of nitroperfluoroacetone with nitrosyl fluoride produces the same perfluoroisopropyl nitrite as that produced by heating of perfluoropropylene with nitrogen dioxide, which proves the structure of nitroperfluoroisopropyl nitrite according to the reaction



which corresponds to structure III. Orig. art. has: 1 table.

SUB CODE: 07/

SUBM DATE: 21Nov63/

ORIG REF: 005/

OTH REF: 005

Card 3/3. BLG

ea FOKIN, A.V.

10

Addition of hydrogen sulfide to mercaptans and olefins
L. Kurnyants and A. V. Fokin. *Ispoln. Khim.* 10
345-44(1950). - Crit. review: 135 references. - N. Thon

1767

Jul/Aug 51

USSR/Chemistry - Plastics

"Polymerization of Fluoroolefins," I.L. Knunyants, A.V. Fokin, Moscow

"Uspekhi Khim" Vol XX, No 4, pp 410-429

Reviews on the basis of published papers the following subjects: tetrafluoroethylene, chlorotri-fluoroethylene (perfluorochlorovinyl), vinylidene fluoride, 1,1-dichloro-2,2-difluoroethylene, vinyl fluoride, haloprenes, vinylfluoroacetates, fluoroacrylates, polyfluorostyrenes. The bulk of the information given is based on foreign publications. Refers to the following Russian work:

191T2

Jul/Aug 51

USSR/Chemistry - Plastics (Contd)

A.P. Borodin's synthesis of organic acid fluorides for the 1st time, A.M. Nesmeyanov's synthesis of formyl fluoride for the 1st time, A.I. Mashen-tsov's method of prepolymer acid fluorides by heating acid chlorides with potassium fluoride ("Zhur Obshch Khim" Vol XV, 1945, p 915).

191T2

(CA 48 no.1:397 '4)

FOKIN, A. V.

191T2

FOKIN, A. V.

Chemical Abst.
Vol. 48 No. 9
May 10, 1954
Organic Chemistry

② Chem
/ Addition reactions of perfluoroblenes. I. L. Knunyants
and A. V. Fokin. *Bull. Acad. Sci. U.S.S.R., Div. Chem.*
Sci. 1952, 279-83 (Engl. translation).—See C.A. 47,
3221b. H.L.H.

Form 4 V.

4

Addition reactions of fluorolefins II: Addition of
alcohols and thiols to perfluoropropylene
A. I. Shchekotikhin and A. V. Golovinskiy
U.S.S.R. Dokl. Akad. Nauk 193, 1937, 1938
See C.A. 48, 1957a.

FORIN, A.V. AND KNUNYANTS, I.I.

Addition Reactions of Perfluoroolefins, Izvestiya, Akademii Nauk SSSR, Otdeleniye
Khimicheskikh Nauk, No. 2, 1952, pp 261-267.

FOKIN, A.V.

112
 Addition reactions of fluorobenzene. II. Addition of alcohols and thiols to perfluoropropylene. J. L. Kamenets, A. I. Shebekotikhin, and A. V. Fokin. *Izv. Akad. Nauk S.S.S.R., Obshch. Khim. Nauk* 1953, 2:2-9; cf. *C.A.*, 47, 3221b. — In the presence of alkali, perfluoropropylene (I) readily adds ROH and RSH with the H atom, adding to the central C atom of I, contrary to the reaction with C_2H_4 . Heating 15 g. I, 3 g. MeOH, and 0.7 g. powd. KOH in a stainless-steel autoclave 13 hrs. at 60° gave 10.1 g. product, b. 44-53°, which, treated with ice cooling with Br until the reaction ceased, and distd., gave 83% pure CF_3CHFCF_2OMe (II), b. 51-5°, d_4^{20} 1.420, n_D^{20} 1.2830; a small amt. of $CF_3BrCF_2CF_2OMe$, b. 136°, d_4^{20} 1.900, n_D^{20} 1.3700, was isolated from the high-boiling residue. Similarly a 14-hr. reaction at 60-60° with EtOH gave 79.3% CF_3CHFCF_2OPr , b. 64-5°, d_4^{20} 1.200, n_D^{20} 1.2060. $PrOH$ gave CF_3CHFCF_2OPr , b. 62-3°, d_4^{20} 1.250, n_D^{20} 1.3110. $iso-PrOH$ after 20 hrs. gave 17.1% CF_3CHFCF_2OCHMe , b. 76°, d_4^{20} 1.280, n_D^{20} 1.3220. $BuOH$ in an 18-hr. reaction gave 34% CF_3CHFCF_2OBu , b. 108°, d_4^{20} 1.270, n_D^{20} 1.3390. $PhOH$ in a 10-hr. reaction gave 44.7% CF_3CHFCF_2OPh , b. 95°, d_4^{20} 1.371, n_D^{20} 1.3965. Heating in a steel autoclave 11.8 g. II, 10 g. concd. H_2SO_4 , and 2.2 g. powd. glass (to bind the HF) 3 hrs. at 60-6° and quenching in H_2O gave 65.6% CF_3CFHCO_2Me , b. 60°, d_4^{20} 1.353, n_D^{20} 1.3192. Similarly was obtained 59.4% *Et ester*, b. 108-0°, d_4^{20} 1.250, n_D^{20} 1.3260. Heating 33 g. I, 10.5 g. MeSH, and 1 g. powd. NaOH in an autoclave 6 hrs. at 120-40° gave 7.5 g. CF_3CHFCF_2SMe , b. 87°, d_4^{20} 1.380, n_D^{20} 1.3443; 12.5 g. EtSH gave 34% (17 g.) CF_3CHFCF_2SEt , b. 100-1°, d_4^{20} 1.322, n_D^{20} 1.3543; distn. of the residue gave an unstated low yield of a disulfide, $C_6F_5S_2H$, b. 80-90°, d_4^{20} 1.291, n_D^{20} 1.4045. The formation of disulfides of the type $RS-CF_2CHFCF_2-SR$ can be explained by the attack of SR ion on I with evolution of HF and formation of allylic $CP_2-CF_2CF_2-SR$, which then adds the 2nd mole of RSH. The formation of allylic by-products is confirmed by the bromination reaction cited above in the ROH series; it is regarded not as direct substitution of F in CP_2 but as a

(over)

4
(2)
2/2 I. L. Kurnyants, A. I. Shchekina, Ushakov
C. A. V. F. Shchekina
result of anionoid attack by the nucleophilic reagent on
the terminal C atom of I having the least electron density,
with the shift of unsatn. and elimination of one F ion, as an
anion. Heating 10 g. $\text{HOCH}_2\text{CH}_2\text{SH}$, 20 g. I, and 0.8 g.
powd. NaOH in an autoclave 6 hrs. at $100-20^\circ$ similarly
gave 20.6% $\text{CF}_3\text{CHFCF}_2\text{SCH}_2\text{CH}_2\text{OH}$, b_p $63-4^\circ$, d₄ 1.546,
n_D²⁰ 1.3835. Heating 10 g. Et_3NH , 20 g. I, and 1 g. borax;
similarly 4 hrs. at 90° treating the mixt. with H_2O gave an
unstated yield of $\text{CF}_3\text{CHFCONEt}_3$, b_p 89° , d₄ 1.228, n_D²⁰
1.3910; presumably the initial product was an unstable
 $\text{CF}_3\text{CHFCF}_2\text{NEt}_3$.
G. M. Kosolapoff

Reactions of chlorides of perfluoro alcohols. II. 412

1,1,1,1-tetrafluoro-2,2,2-trifluoroethane (1,1,1,1-tetrafluoro-2,2,2-trifluoroethane) was treated with $\text{CF}_3\text{CF}_2\text{SH}$ with cyclohexane similarly gave an unstated yield

F. 11/14/55
KNUNYANTS, I.L., akademik; FOKIN, A.V.

Fluorine and its compounds. Priroda 44 no.8:3-19 Ag '55.
(Fluorine) (MIRA 8:10)

Translation D 492394

FOKIN, A.V.

USSR/Organic Chemistry. Synthetic Organic Chemistry.

G-2

Abs Jour: Referat Zhur-Khimiya, No 4, 1958, 11392.

Author : Khunyanys, I. L. and Fokin, A. V.

Inst : Academy of Sciences USSR

Title : The Nitration of Perfluoroolefins by Nitrogen Dioxide

Orig Pub: Doklady Akad Nauk SSSR, 111, No 5, 1035-1038 (1956)

Abstract: The reaction of N_2O_4 with perfluoroolefins proceeds by a free radical mechanism and leads to the formation of dinitroperfluoroalkanes and β -nitroperfluoroalkylnitrites; the overall yield is 90%. The reactivity of the perfluoroolefins decreases from left to right in the following series $CF_2=CF_2$ (I) $>$ $CF_3CF=CF_2$ (II) $>$ $CF_2CF=CF_2$ (III) $>$ $(CF_3)_2C=CF_2$ (IV). I in CCl_4 , $CHCl_3$, CCl_2F_2 , or $CClF_2CClF_2$ at 20° (in the absence of a solvent the reaction proceeds explosively)

Card : 1/3

USSR/Organic Chemistry. Synthetic Organic Chemistry.

G-2

Abs Jour: Referat Zhur-Khimiya, No 4, 1958, 11392.

gives equal amounts of $CF_2(NO_2)CF_2NO_2$ (throughout this abstract the characteristics of the products are given in the order bp in °C, n_D^{20} , and d_4^{20}), 57-58, 1.3249, 1.622 and $CF_2(NO_2)CF_2ONO$, 17, 1.3002 (0°), 1.5310 (0°). The nitration of II at 100° in the absence of a solvent gives $CF_3CF(NO_2)CF_2NO_2$ (in insignificant amounts), 76, 1.3141, 1.6313 and, as the main product, $CF_3CF(NO_2)CF_2ONO$ (V), 57, 1.5276, 1.637. The reaction of III with N_2O_4 proceeds only in a narrow temperature range (130-160°) and leads to the formation of $CF_2CF(NO_2)CF(NO_2)CF_2$, 62°/100 mm, 1.3640, 1.7721, and $CF_2CF(ONO)CF(ONO)CF_2$, 20, -, 1.5481. IV reacts with N_2O_4 at 180° giving (equal amounts) $(CF_3)_2C(NO_2)CF_2NO_2$, 92, 1.3212, 1.660, and $(CF_3)_2C(NO_2)-CFONO$ (VI), 48, 1.2870, 1.598. The structure

Card : 2/3

AUTHORS:

Fokin, A.V.
Knunyants, I.L., Fokin, A.V.

62-12-3/20

TITLE:

Nitration of Fluorine Olefines by Nitrogen Dioxide (Nitrovaniye fluorolefinov dvuckis'yu azota). Lecture Delivered at the Meeting of the Department of Chemical Sciences AN USSR on October 30, 1957 (Doklad na sessii Otdeleniya khimicheskikh nauk Akademii nauk SSSR 30 oktyabrya 1957 g).

PERIODICAL:

Izvestiya AN SSSR Otdeleniya Khimicheskikh Nauk, 1957, Nr 12, pp. 1439-1451 (USSR)

ABSTRACT:

The reactions of the nitration of saturated and unsaturated hydrocarbons, which have already been thoroughly investigated, were practically not investigated at all in fluorine-organic compounds. The majority of the reaction of fluorine olefines with "nucleophylic" reagents has ionic character. In contrast to the smooth interaction of perfluorine-olefines with nucleophylic reagents, their interaction with electrophylic substances takes place under much more difficult conditions. According to Koffman (1949) and Khastsel'din (1953) only dinitroalkanes are formed by the interaction of tetrafluorine ethylene and chlorine trifluorineethylene with nitrogen dioxide. Reactions with other fluorine olefines were not investigated. Among other things, the authors found that these reactions

Card 1/3

Nitration of Fluorine Olefines by Nitrogen Dioxide.
Lecture Delivered at the Meeting of the Department
of Chemical Sciences AN USSR on October 30, 1957

62-12-3/20

depend mainly on the structure of the characteristic features of fluorine olefine (and lead to the formation of new and interesting substances). It was further shown that tetrafluorine-ethylene reacts explosively with nitrogen dioxide. It was possible to extend the method of nitration by means of nitrogen-dioxide also to other perfluorine olefines (see tables). In the case of none of the methods investigated were compounds able to form. It was shown that the destruction of β -nitroperfluorine-ethyl and β -nitroperfluorpropylnitrites begins only at a temperature of more than 250°C . The investigation of the nitration of chlorine fluorine olefines made it possible to determine a certain characteristic feature of this reaction (see formulae on page 1446.) The investigation of the nitration reactions of fluorine olefines and not substituted olefines with nitrogen-dioxide made it apparent that there is a similarity of the chemical character of these reactions (see table 2). The results of this investigation further showed that the stability of intermediate radicals as well as the polarity of fluorine olefines and that of the radical-like particle NO_2 are an important factor

Card 2/3

Nitration of Fluorine Olefines by Nitrogen Dioxide.
Lecture Delivered at the Meeting of the Department
of Chemical Sciences AN USSR on October 30, 1957

62-12-3/20

of orientation of the reacting components. Conceptions concerning the polarity of radicals, which were first published by Uoters (Waters?) and were further developed by Karash, Veys, Dolgopolo and others, deserve attention. There are 2 tables and 18 references, 10 of which are Slavic.

SUBMITTED: October 9, 1957

AVAILABLE: Library of Congress

Card 3/3

1. Chemical engineering-Conference
2. Hydrocarbon-Reactions
3. Fluorine-Organic compounds
4. Fluorine olefines

1 FOKIN, A.V.

Nitroperfluorobutyl nitrate
 b. 118°, 1.638, 1.3760; $(CF_3)_2C(NO_2)CO_2H$ b. 117.5°, 1.412, 1.3470; $(CF_3)_2C(NO_2)CO_2Et$ b. 53°, 1.3790, 1.3567; O_2NCF_2COCl b. 52°, 1.3581; $(O_2NCF_2CO)_2O$ b. 83°, 1.2820, 1.3596; $(CF_3)_2C(NO_2)CN$ b. 27°, 1.4010, 1.3220; $(CF_3)_2C(NO_2)CO_2CH_3$ b. 134°, 1.441, 1.3362; $CF_3CF(NO_2)COCl$ b. 37°, 1.6590, 1.3502; $CF_3CF(NO_2)CONEt_2$ b. 149°, 1.398; $O_2N(CF_2)_2NO_2$ b. 110°, 1.703, 1.3410; $H_2N(CF_2)_2CONH_2$ --, -- (m. 250°). Dissociation constant of $CF_3CF(NO_2)CO_2H$ is $0.7,000 \times 10^{-4}$; that of $CF_3CF(NO_2)CO_2H$ is 1.875×10^{-4} . The metallic salts of the acids are unstable. II Ag salt decomp. to the salt of perfluoropropionic acid. Attempts to prep. $CF_3CF(NO_2)CN$ gave instead $CF_3CF(NO_2)CO_2H$. Nitration of perfluorocyclobutane yields an acid which produces some perfluorocyclobutanediol 1,2-dinitro compound (C.A. 51, 9472n), which hydrolyzes to perfluorocyclobutanediol; treatment with NH_3 yields the amide; treatment with alic. yields esters. G. M. K.

FOKIN, A. V.

"Prospects of Using Fission Product Source Radiation in Radiation Chemistry",

by N. V. Zimakov, E. V. Volkova, A. V. Fokin, V. V. Kulichenko, V. G. Vereskunov,
A. G. Bykov, and N. I. Bogdanov

Report presented at 2nd UN Atoms-for-Peace Conference, Geneva, 9-13 Sept 1958

FOKIN, A-V

AUTHOR: None given

62-58-4-30/32

TITLE: **Anniversary Session of the Department**
for Chemical Sciences of the AS USSR on October 30 and 31,
1957, and General Meeting of the Department for Chemical
Sciences on December 19 and 20, 1957 (Yubileynaya sessiya
otdeleniya khimicheskikh nauk Akademii nauk SSSR ot 30-31
oktyabrya 1957 g. i obshcheye sobraniye otdeleniya khimicheskikh nauk 19-20 dekabrya 1957 g.)

PERIODICAL: Izvestiya Akademii Nauk SSSR, Otdeleniye Khimicheskikh Nauk,
1958, Nr 4, pp. 521 - 524 (USSR)

ABSTRACT: On the occasion of the 40th anniversary of the October
Revolution a reunion meeting of the Department for Chemical
Sciences of the AS USSR took place. In his opening speech
N. N. Semenov pointed out the outstanding success of the USSR
in the field of sciences especially in that of chemistry.
Scientific lectures of the sessions were held by the following
scientists, as was mentioned already earlier: Knunyants,
Member, Academy of Sciences, and A. V. Fokin on the "Nitration
of Fluorofines", A. L. Midzhoyan, Member, AS Armenian SSR,

Card 1/4

62-58-4-30/32

Anniversary Session of the Department for Chemical Sciences of the AS USSR on October 30 and 31, 1957, and General Meeting of the Department for Chemical Sciences on December 19 and 20, 1957.

on the "Investigations in the Field of the Synthesis of Physiologically Active Compounds", R. Kh. Freydlina, Doctor of Chemical Sciences, reported on the "Investigation of the Telomerization Reaction and the Reaction of the Synthesis on the Basis of Telomers" (Reference 2). B. A. Dolgoplosk, Doctor of Chemical Sciences, spoke on the "Generation of Free Radicals in Solutions and Their Reactions in Model Systems," A. M. Frumkin, Member of the Academy of Sciences, reported on "Some General Problems of Electrochemical Kinetics and the Theory of Ion Reactions" (Reference 4), A. V. Kiselev, Doctor of Chemical Sciences (Reference 5) spoke on "Some Problems of Adsorption Theory", N. M. Emanuel' (Reference 6), Doctor of Chemical Sciences, reported on "New Problems in the Field of Chain Reactions", V. L. Tal'roze, Candidate of Chemical Sciences, spoke on mass-spectroscopic investigations of ion-and radical reactions, A. P. Rebinder, Member,

Card 2/4

62-58-4-30/32

Anniversary Session of the Department for Chemical Sciences of the AS USSR on October 30 and 31, 1957, and General Meeting of the Department for Chemical Sciences On December 19 and 20, 1957

Academy of Sciences, drew conclusions with regard to the development of physico-chemical mechanics (Reference 7). I. V. Tananayev, Corresponding Member of the AS USSR, gave new data on the chemistry of some rare elements, D. I. Ryabchikov and others spoke on the "Problems of the Chemistry of Rare Earth Elements"; the final lecture was that of V. A. Sokolov, Doctor of Chemical Sciences, on the "Calorimetric Measurements at High Temperatures". General Regular Meeting of the Department for Chemistry of the AS USSR (December 19 - 20, 1957): A. I. Brodskiy, Corresponding Member, AS USSR, spoke on the "Investigation of Some Reactions of Peroxides and Peracids of Hydrogen by Means of the Isotopic Method", M. M. Shemyakin, Corresponding Member, AS USSR, spoke on the "Use of N^{15} for the Explanation of the Mechanism of Some Organic Reactions", O. A. Reutov, Doctor of Chemical Sciences, reported on the "Investigation of the

Card 3/4

62-58-4-30/32

Anniversary Session of the Department for Chemical Sciences of the AS USSR on October 30 and 31, 1957, and General Meeting of the Department for Chemical Sciences On December 19 and 20, 1957

Electrophil and Homolytical Reactions of the Substitution in the Carbon Atom by Means of the Method of Isotope Exchange", I. P. Alimarin, Corresponding Member, AS USSR, reported on new methods of determination of the division of rare elements using organic derivative sulfuric-, selenic- and telluric acids, V. G. Levich, Doctor of Chemical Sciences, reported on the "Diffusion Kinetics of Heterogenous Chemical Reactions in mobile Liquids". There are 8 references, all of which are Soviet.

AVAILABLE: Library of Congress

1. Chemical Industry--USSR

Card 4/4

FOKIN, A.V.; VOLKOVA, Ye.V.; SOROKIN, A.D.

Utilization of energy of ionizing radiations in the process of polymerization of trifluoroethylene. Polymerization of trifluoroethylene in block and in the medium of chlorine-containing solvents. Khim.nauka i prom. 4 no.6:806-807 '59. (MIRA 13:8)

(Ethylene)
(Polymerisation)
(Gamma rays)

81139

S/064/60/000/03/02/022
B010/B008

5.3600

AUTHORS: Fokin, A. V., Doctor of Chemical Sciences,
Kosyrev, Yu. M., Candidate of Technical Sciences

TITLE: Pyrolysis of Carbon Fluorides

PERIODICAL: Khimicheskaya promyshlennost', 1960, No. 3, pp. 186-192

TEXT: The pyrolysis of poly-tetrafluoroethylene at atmospheric pressure as well as in vacuum (1 torr) was studied in an apparatus (Figs. 1,2) designed for the pyrolysis of solid substances. Diagrams (Figs. 3,4) illustrate the change of the composition of the pyrolyzate with the pyrolysis temperature. The composition of the pyrolyzate obtained at atmospheric pressure is given in Table 1 and that of the product obtained in vacuum in Table 2. The range of 600-750°C proved to be most favorable for atmospheric pressure, a maximum yield of perfluorocyclobutane (>57%) being obtained at 600°C, of perfluoropropylene (45%) at 700-710°C, and of perfluoroisobutylene (33%) at 750°C. The pyrolysis of perfluorocyclobutane was studied in the steel tube (steel of the grade NT5Kh92Z6 (NT5Kh92Z6)) of a device used for the pyrolysis of gaseous carbon fluorides at contact

Card 1/2

81139

Pyrolysis of Carbon Fluorides

S/064/60/000/03/02/022
B010/B008

times of 30 sec. and temperatures of 650-800°C. It was established that this pyrolysis is very similar to that of poly-tetrafluoroethylene, and that perfluorocyclobutane can thus also be used for the production of perfluoropropylene and perfluoroisobutylene by pyrolysis. Studies of the pyrolysis of tetrafluoroethylene showed analogous temperature dependences of the process and, thus, also of the reaction mechanism of the pyrolysis of perfluorocyclobutane and poly-tetrafluoroisobutylene (Table 3). Further experiments showed that the production of perfluoroisobutylene by pyrolysis of perfluoropropylene takes place best at a contact duration of 70 sec. and a temperature of 700-710°C (Fig. 7). Experiments on perfluoroisobutylene pyrolysis showed (Table 4) that the latter is the most heat-resistant fluoro olefine. The transformations of the compounds under consideration are schematically shown in Fig. 8 on the basis of the experimental results obtained, and the authors point out that the remains of poly-tetrafluoroethylene production can be used for the production of perfluoropropylene and perfluoroisobutylene. There are 8 figures, 4 tables, and 8 references: 3 Soviet, 1 Canadian, 1 British, and 2 American.

Card 2/2

82356

S/063/60/005/001/009/009

5.2831

AUTHORS: Fokin, A. V., Volkova, Ye. V., Sorokin, A. D.

TITLE: On the Use of the Energy of Ionizing Radiation in the Process of Copolymerization of Trifluorochloroethylene With Various Monomers

PERIODICAL: Zhurnal vsesoyuznogo khimicheskogo obshchestva im. D. I. Mendeleeva, 1960, Vol. 5, No. 1, p. 120

TEXT: The possibility was shown of radiation copolymerization¹⁹ of trifluorochloroethylene with various perfluorinated and partially fluorinated olefines and also with ethylene oxide. Vinylidenefluoride, perfluoropropylene, tetrafluoroethylene and ethylene oxide were used as second components in the copolymerization under the action of γ -radiation. The experiments were carried out at room temperature in metal ampoules made of β Q-1T (EYa-1T) stainless steel. The copolymerization of trifluorochloroethylene with vinylidenefluoride was carried out in the molar ratio $CF_2 = CFCI : CH_2 = CF_2$ from 3 : 1 to 1 : 3 at a dose intensity of 14-16 r/sec and a dose of 2-3 million r. Under these conditions practically the complete conversion of both monomers is obtained. The radiation-chemical yield is 3-5,000 molecules per 100 ev. The copolymer obtained is sufficiently resistant against alcohols, various oils and nitric acid; it is

Card 1/2

82356

S/063/60/005/001/009/009/

On the Use of the Energy of Ionizing Radiation in the Process of Copolymerization of Trifluorochloroethylene With Various Monomers

soluble in diethyl ether, acetone and esters. The copolymers of trifluorochloroethylene with perfluoropropylene, trifluorochloroethylene with tetrafluoroethylene and the polymer of vinylidene fluoride were obtained under analogous conditions. A copolymer of trifluorochloroethylene with ethylene oxide was obtained under the action of γ -radiation of Co^{60} . There are 3 tables and 3 references: 2 Soviet and 1 American. ✓

SUBMITTED: September 30, 1959.

Card 2/2

PETROV, K.A.; SHEVCHENKO, V.B.; TIMOSHEV, V.G.; MAKLYAYEV, F.A.; FOKIN,
A.V.; RODIONOV, A.V.; BALANDINA, V.V.; YEL'KINA, A.V.; MAGNIBEDA,
Z.I.; VOLKOVA, A.A.

Alkyl phosphonates, diphosphonates, and phosphine oxides as
extracting agents. Zhur.neorg.khim. 5 no.2:498-502

F '60. (MIRA 13:6)

(Phosphonic acid) (Phosphine oxide)

(Extraction(Chemistry))

35437

S/081/62/000/004/074/087
B138/B110

112214

AUTHORS: Zimakov, P. V., Volkova, Ye. V., Fokin, A. V., Sorokin, A. D.,
Belikov, V. M.

TITLE: Use of nuclear radiation energy in the process of the
polymerization of fluoro-olefines

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 4, 1962, 557, abstract
4P24 (Sb. "Radioakt. izotopy i yadern. izlucheniya v nar.
kh-ve SSSR, v. 1. M.", Gostoptekhizdat, 1961, 219-226)

TEXT: The processes of the separate and combined radiation polymerization
of tetrafluorethylene and trifluorchlorethylene have been investigated with
the aim of eliminating some of the deficiencies in existing methods of
fluoro-olefine polymerization. It has been found that tetrafluorethylene
and trifluorchlorethylene can easily be polymerized under various
temperature conditions and mediums with comparatively low radiation
intensities. The resulting polymers have a high degree of purity. The
possibility of producing various fluoro-copolymers by radiation is
demonstrated. Both radiation polymerization and radiation vulcanization
might be carried out in the case of fluor-containing rubbers. [Abstracter's
note: Complete translation.]
Card 1/1

VOLKOVA, Ye.V.; FOKIN, A.V.; BELIKOV, V.M.

Polymerization of tetrafluoroethylene by the action of gamma
rays. Zhur.VKHO 6 no.1:113-114 '61. (MIRA 14:3)
(Ethylene) (Gamma rays) (Polymerization)

FOKIN, A.V.; SKLADNEV, A.A.; KNUNYANTS, I.L., akademik

Reactions of fluorinated olefines. Reactions between fluorinated
olefins and hydrogen sulfide. Dokl.AN SSSR 138 no.5:1132-1135 Je
'61. (MIRA 14:6)

(Olefins)

(Hydrogen sulfide)

S/844/62/000/000/078/129
D423/D307

AUTHORS: Volkova, Ye. V., Fokin, A. V. and Sorokin, A. D.

TITLE: Radiation polymerization of trifluorochlorethylene

SOURCE: Trudy II Vsesoyuznogo soveshchaniya po radiatsionnoy khimii. Ed. by L. S. Polak. Moscow, Izd-vo AN SSSR, 1962, 460-464

TEXT: The present work was carried out in view of the inconclusive results obtained by other workers on the radiation polymerization of F-containing unsaturated organic compounds. Experiments were carried out in stainless steel flasks. It was not possible to determine the initial velocity of polymerization but the relationship between velocity, temperature and radiation dosage was determined over linear sections of the kinetic curves corresponding to 5 - 20% conversion. It was established that polymerization takes place over a wide range of dosage and that it proceeds without a significant induction period. The power index of the velocity-dosage relationship varied from 0.36 to 0.5 for higher dosages, and the relationship was charac-

Card 1/2

Radiation polymerization of ...

S/844/62/000/000/078/129
D423/D307

teristic for chain reaction processes. The presence also of a radiation 'after-effect' was shown, which lasted over a period of 2 days. Between -21 and $+60^{\circ}\text{C}$ the rate of polymerization increased initially to a maximum at 35°C and then fell off, confirming the results obtained by Roberts. Over the temperature range studied, the radiation yield amounted to $27,000$ mol/100 ev absorbed energy for a dose of 3×10^5 rad/hr. On increasing the temperature from 0°C to 35°C , a reduction in the molecular weight of the polymer was observed and this value was also reduced at higher doses. The experiments indicated that radiation polymerization of trifluorochlorethylene takes place by a chain process, originated by a radical mechanism. There are 5 figures.

Card 2/2

S/844/62/000/000/079/129
D423/D307

AUTHORS: Volkova, Ye. V., Fokin, A. V., Zimakov, P. V. and Belikov, V. M.

TITLE: Certain special features of the radiation polymerization of tetrafluorethylene by the action of β and γ radiations

SOURCE: Trudy II Vsesoyuznogo soveshchaniya po radiatsionnoy khimii. Ed. by L. S. Polak. Moscow, Izd-vo AN SSSR, 1962, 465-469

TEXT: Recent investigations are described of the radiation polymerization of TFE in the solid, liquid and vapor phases, using Co^{60} and Sr^{90} as the γ and β sources. Irradiation in the liquid phase was carried out at 20 - 25°C using CHCl_3 as the solvent with a dose-rate of 11 rad/sec. Conversion of monomer increased with increase of dosage and concentration of monomer. The polymer obtained (PTFE) contained up to 2% chlorine, which was explained by the fact that the CHCl_3 also participates in the reaction by interaction of

Card 1/2

Certain special features ...

S/844/62/000/000/079/129
D423/D307

radicals and chain breakage in the polymer, resulting in low-molecular weight PTFE. Experiments in the solid state were carried out from -80 to 0°C with dosages of 1×10^5 and 1×10^4 rad. Almost total conversion of monomer occurred after 200 min at -80°C and after 20 mins at 0°C. Exceptionally large yields were obtained in comparison with similar reactions of other unsaturated compounds. The existence of a radiation after-effect was confirmed, which continued over several hours after removal of the radiation source. Experiments in the gas phase showed the presence of an induction phase extending over several hours. After the appearance of solid PTFE the reaction velocity was increased. The temperature was maintained at 20 - 25°C and a Sr^{90} B source was used with a dose-rate of 5 rad/sec. Results indicated a high tendency of TFE towards radiation polymerization with a high yield (approx. 10^6 mol/100 ev absorbed). There are 5 figures and 1 table.

Card 2/2

11119

S/063/62/007/005/006/006

A057/A126

11 22 14
AUTHORS: Volkova, Ye.V., Fokin, A.V., Sorokin, A.D., Bulygina, L.A.

TITLE: On the polymerization of vinylidenfluoride under the influence of γ -irradiation

PERIODICAL: Zhurnal vsesoyuznogo khimicheskogo obshchestva imeni D.I. Mendeleeva, v. 7, no. 5, 1962, 593 - 594

TEXT: Radiative polymerization "in bulk" of vinylidenfluoride was investigated and the obtained results compared with previous studies carried out with tetrafluoroethylene and trifluorochlorethylene. The rate of radiative polymerization under same conditions lies in the sequence tetrafluoroethylene > vinylidenfluoride > trifluorochlorethylene and the corresponding yields per 100 ev are 10^6 , 10^5 , and 10^4 molecules, respectively. The polymerization occurs in all cases with a high conversion rate, practically up to 100%. The present experiments were made in 25 ml 1X18H9T (1Kh18N9T) steel autoclave test tubes, using a Co^{60} source with a total capacity of 5,000 g.equiv. Ra. A considerable induction period, effected by impurities (especially oxygen), was observed and, therefore, the monomer purified before use. The latter was a commercial grade of 99.8% purity. The de-

Card 1/2

On the polymerization of.....

S/063/62/007/005/006/006
A057/A126

pendence of the conversion upon the irradiation time was studied at 23°C with doses of 1, 5, 10, and 30 rad/sec and the rate of reaction determined from the inclination of the kinetic curves. The value of the radiation-chemical yield decreases with the dose capacity. An increase in temperature raises the rate of the radiation polymerization but for all investigated temperatures (-78, -20, 0, 23°C at 10 rad/sec) a maximum value was obtained after about 6 h. The total activation energy of radiative vinylidenefluoride polymerization was determined with 3.6 kcal/mole. The process occurs by a radical-chain mechanism.

SUBMITTED: May 12, 1962

Card 2/2

KNUNYANTS, I.L.; FOKIN, A.V.; KOMAROV, V.A.

Nitration of perfluoropropylene with nitrogen dioxide.

Zhur. VKHO 7 no.6:709-710 '62.

(MIRA 15:12)

(Propene)

(Nitrogen oxide)

FOKIN, A.V.; SKLADNEV, A.A.; STUDNEV, Yu.N.; KNUNYANTS, I.L., akademik

Interaction of asymmetric fluoroolefins with hydrogen sulfide.
Dokl. AN SSSR 142 no.1:99-101 Ja '62. (MIRA 14:12)
(Olefins) (Hydrogen sulfide)

KNUNYANTS , I.L., akademik; FOKIN, A.V.; BLAGOVESHCHENSKIY, V.S.; KOSYREV, Yu.M.

New interesting cases of the formation of nitroso compounds.

Dokl. AN SSSR 146 no.5:1088-1091 0 '62.

(MIRA 15:10)

(Nitroso compounds)

FARKH, A. V.

PHASE I BOOK EXPLOITATION

SOV/6488

Knunyants, Ivan Lyudvigovich, Academician, and Aleksandr
Vasil'yevich Fokin, Professor

Pokoreniye nepristupnogo elementa (Conquest of an Inaccessible
Element) Moscow, Izd-vo AN SSSR, 1963. 189 p. (Series:
Akademiya nauk SSSR. Nauchno-populyarnaya seriya) Errata
printed on inside of back cover. 25,000 copies printed.

Ed. of Publishing House: V. M. Tarasenko; Tech. Ed.: S. P.
Golub'.

PURPOSE: This textbook is intended for chemists and engineers.

COVERAGE: The book covers the full range of fluorine chemistry
and technology and is based on Soviet and Western sources. The
text includes: properties of fluorocarbons; fluorinated
hydrocarbons, e.g., fluorinated olefins which have high
thermal stability and chemical resistance; esters of dibasic

Card 1/³₈

Conquest of an Inaccessible Element

SOV/6488

perfluorocarboxylic acids which are used in the preparation of stable lubricants and in synthetic rubber; preparation, properties, and applications of fluorine-containing plastics [teflon, teflon-100, poly(vinylidene fluoride)]; their radiation polymerization; fluorine containing elastomers used in supersonic aircraft and jet engines; freons; and other applications. The fifth chapter is devoted to lubricants and hydraulic fluids with high thermal and oxidation stability, which can be used under severe conditions and in systems with aggressive media. It is noted that chlorofluorocarbon oils lower the friction coefficient of rubbing surfaces made of the same metal under heavy loads and at high temperatures. Certain Soviet oils, greases, and hydraulic fluids are also described. Tabulated data are given for: 4-F, 3-F, 3-OK, 10-OK, 20-F, Summer No.5, Winter No.8, UPI, Fluid No.12, manometric fluid, and balancing fluid. Only KS, UPI, and Summer No.5 are insoluble in ammonia and amines. In describing esters of dicarboxylic acids and fluorinated

Card 2/5

Conquest of an Inaccessible Element

SOV/6488

alcohols, the author mentions the applicability of such lubricants for submarines to avoid a detectable oil film on the water surface (the ONR research is quoted). Modern lubricants are already in use in jet engines and other engines working at high temperatures. The author points out the importance of fluorine compounds in rocketry, aviation, astronautics, atomic energy, and industry. The last chapter is devoted to fluorine organic compounds used for: the improvement of light fastness of dyes (benzotri-fluoride, fluorobenzene, trifluoromethylbenzene); the preparation of chemotherapeutic compounds; and the preparation of a class of toxic compounds (fluoroacetates). These compounds belong to esters of the general type $F(Ch)_n COOR$, fluoroalcohols, and fluoroacetamides with their derivatives. Fluorophosphates are mentioned as CW compounds (e.g., DFP-3, Sarin, Soman) which caused a change in anti-CW methods. No personalities are mentioned. There are no references.

Card 3,³₅

KNUMYANTS, I. L.; FOKIN, A. V.; DYATKIN, B. L.; KOMAROV, V. A.

Action of nitrogen dioxide on perfluoroisobutylene. Zhur.
VKHO 8 no.2:239-240 '63. (MIRA 16:4)

1. Institut elementoorganicheskikh soyedineniy AN SSSR.

(Nitrogen oxides) (Propene)

FOKIN, A.V.; STUDNEV, Yu.N.; SKLADNEV, A.A.

Reactions of 1,1-difluoropolyfluoralkylmercaptans with phosphorus acid derivatives. Zhur.ob.khim. 33 no.10:3366-3369
0 '63.
(MIRA 16:11)

FOKIN, A.V.; SKLADNEV, A.A.; KOMAROV, V.A.

Acylating action of mixed anhydrides of fluorine-containing
carboxylic acids. Zhur.ob.khim. 33 no.10:3271-3274 0 '63.
(MIRA 16:11)

L 53233-65 EWT(m)/EPF(c)/EPR/EWP(j)/EWA(c)
ACCESSION NR: AP6016225 JW/RM

Pc-4/Pr-4/Ps-1 RPL WW/
UR/0063/65/010/003/0354/0355
542.958.1 + 547.32


AUTHOR: Fokin, A.V.; Komarov, V.A.; Borochkin, I.N.; Davydova, S.M.

TITLE: Nitration of 1,1,1-trifluoropropylene by nitrogen dioxide and a study of the nitration products

SOURCE: Vsesoyuznoye khimicheskoye obshchestvo. Zhurnal, v. 10, no. 3, 1965, 354-355

TOPIC TAGS: nitration, olefin, nitrogen oxide, nitration product

ABSTRACT: The nitration of olefins having the general formula $Rf-CH=CH_2$ (where $Rf = CF_3, C_3F_7-CH_2-CF_2$) was studied. Because the individual compounds cannot be separated by ordinary fractionation of the reaction mixture, the reaction products were treated with water, the reaction mixture was extracted with ethyl ether, and the ether solution was dried and fractionated. 3-Nitro-1,1,1-trifluoro-2-propanol (I, 80% yield) and nitro-1,1,1-trifluoro-2-propanol nitrate (II, 10% yield) were obtained. Compound II was also obtained by treating I with a nitrating mixture:




Card 1/3


L 53937-65

ACCESSION NR: AP5018225


Bromination of I in an alkaline medium produced 3-nitro-3-bromo-1,1,1-trifluoro-2-propanol:



and the reaction of I with acetyl chloride yielded 1,1,1-trifluoro-3-nitro-2-propanol:




On the basis of the products obtained, the nitration of 1,1,1-trifluoropropylene may be represented as follows:



Card 2/3

52037-65
ACCESSION NR: AP5016225



Orig. art. has: 5 formulas.

ASSOCIATION: none

SUBMITTED: 28Aug64

ENCL: 00

SUB CODE: OC

NO REF SOV: 000

OTHER: 004

Card 3/3

L 35524-65 EWT(m)/EPF(c)/EPR/ENP(j)/ENP(t)/ENP(b)/ENP(s) 10-4/1-4/10-4 13P(c)/
RPL JD/WW/RM

ACCESSION NR: AP5008203

S/0286/65/000/005/0071/0071

AUTHORS: Pokin, A. V.; Skladnev, A. A.; Kvashan, Z. N.; Studnev, Y. N. 35

TITLE: A method for producing sulfur-bearing polyfluororganic compounds. Class 19,
No. 168882

Source: 'Byulleten' izobreteniy i tovarnykh znakov, no. 5, 1965, 71

TOPIC TAGS: sulfur, fluorine, organic derivative, olefin

ABSTRACT: This Author Certificate presents a method for obtaining sulfur-bearing polyfluororganic compounds. To expand the raw-material base, fluoroolefin is made to react with hydrogen sulfide in the presence of an alkali metal salt.

REMARKS: none

SUBMITTED: 12Dec61

ENCL: 00

SUB CODE: 00

NC REF SOV: 000

OTHER: 000

Card 1/1

LIMAR', T.F.; UVAROVA, K.A.; BULACHEVA, A.F.; SGYVUBM, A.S.; BEDNOVA, I.N.; MAKOVSKAYA, E.B.; SOLOMEINA, G.I.; DOLMATOV, Yu.D.; BOBYPENKO, Yu. Ya.; KOGAN, F.I.; KOVALENKO, P.N.; IVANOVA, Z.I.; FOKIN, A.V.; KOMAROV, V.A.; SOROCHKIN, I.N.; DAVYDOVA, S.M.; RAVDEL', A.A.; GORELIK, G.N.; DAUKSHAS, V.K. [Dauksas, V.]; PIKUNAYTE, L.A. [Pikunaite, L.]; SHARIPOV, A.Kh.; SHABALIN, I.I.; STEPNOVA, G.M.; SHMIDT, Ye.V.; DUBOV, S.S.; STRUKOV, O.G.

Scientific research papers of the members of the All-Union Mendeleev Chemical Society (brief information). Zhur. VHKO 10 no.3:350-360 '65. (MIRA 18;8)

1. Donetskii filial Vsesoyuznogo nauchno-issledovatel'skogo instituta khimicheskikh reaktivov i osobo chistykh khimicheskikh veshchestv (for Limar', Uvarova, Bulacheva). 2. Ural'skiy nauchno-issledovatel'skiy khimicheskii institut (for Shubin, Bednova, Makovskaya, Solomeina). 3. Chelyabinskiy filial Gosudarstvennogo nauchno-issledovatel'skogo i proyektного instituta mineral'nykh pigmentov (Dolmatov, Bobyrenko). 4. Rostovskiy-na-Donu universitet (for Kogan, Kovalenko, Ivanova). 5. Leningradskiy tekhnologicheskii institut imeni Lensoveta i Institut mineral'nykh pigmentov (for Ravdel', Gorelik). 6. Vil'nyusskiy gosudarstvennyy universitet imeni Kpsukasa (for Daukshas, Pikunayte). Nauchno-issledovatel'skiy institut neftekhimicheskikh proizvodstv (for Sharpipov, Shabalin). 8. Tomskiy politekhnicheskii institut imeni Kirova (for Stepnova, Shmidt).

L 18545-66

ACC NR: AP6002180

(N)

SOURCE CODE: UR/0146/65/008/006/0108/0113

AUTHOR: Vinogradov, A. A.; Fokin, A. V.

ORG: Odessa Higher Marine-Engineering School (Odesskoye vyssheye inzhenernoye morskoye uchilishche)

TITLE: Centrifugal tangential angular-velocity sensor

SOURCE: IVUZ. Priborostroyeniye, v. 8, no. 6, 1965, 108-113

TOPIC TAGS: sensor, angular velocity sensor, automatic control

ABSTRACT: The development is reported of a new sensor (see figure below) which measures simultaneously both the angular velocity and the angular acceleration. Essentially, this is the classical Watt regulator in which axis $O_3 O_4$ of suspension of weight 1 is turned, together with the weight, about axis $O_5 O_6$ (which is parallel to $O_1 O_2$) by an angle θ ; this design modification permits measuring both the velocity and the acceleration. Differential equations for the dynamics of weight 1 and disk 3 are set up and transformed which results in a single second-order differential equation describing the motion of the entire centrifugal tangential sensor. The sensor is intended for turboprop, turbojet, and diesel engines. Orig. art. has: 1 figure, 30 formulas, and 1 table.


SUB CODE: 13 / SUBM DATE: 11Jul64 / ORIG REF: 005

Card 1/2

UDC: 62-552

L 18545-66

ACC NR: AP6002180



Card 2/2 mgs

L 16471-66 EWT(m)/ETC(f)/EPF(n)-2/EWG(m)/EWP(j) WW/DM/RM
ACC NR: AP6005532 (A) SOURCE CODE: UR/0089/66/020/001/0053/0054

AUTHOR: Fokin, A. V.; Kuzicheva, V. S.; Fomin, Yu. K.

43
40
8

ORG: none

TITLE: Possibilities of "oil" flotation for reprocessing liquid radioactive wastes

SOURCE: Atomnaya energiya, v. 20, no. 1, 1966, 53-54

19,55

TOPIC TAGS: flotation, radioactive waste disposal, radioisotope, nuclear engineering, solvent extraction

ABSTRACT: "Oil" flotation may be used at ordinary temperatures with comparatively simple equipment for extracting the solid phase from waste radioactive pulp and concentrating it together with trapped radioisotopes in a layer of organic matter which is immiscible with water. The suspended particles are treated with various water-repellent surface-active sorbents, (e. g. salts of fatty acids). Up to 90-95% of the radioactive isotopes may be removed from the water in a single stage. It is recommended that nonflammable and low-boiling solvents of the carbon tetrachloride type should be used in quantities of 30-50 ml per gram of solid residue to

Card 1/2

UDC: 621.039.722 + 621.928.5

2

L 16471-66
ACC NR: AP6005532

be extracted. In some cases organic monomers may be used for the "oil", and the layer of extracted material may be directly converted to a solid plastic by bulk or suspension polymerization. It was found that preparations based on polystyrene and various polyester acids may be used for burial of the radioactive isotopes.

SUB CODE: 18/ SUBM DATE: 15Oct65/ ORIG REF: 000/ OTH REF: 000

Card 2/2mc

ACC NR: AT6034055

(A)

SOURCE CODE: UR/0000/66/000/000/0109/0114

AUTHOR: Volkova, Ye. V.; Zimakov, P. V.; Fokin, A. V.; Sorokin, A. D.; Belikov, V. M.; Bulygian, L. A.; Skobina, A. I.; Krasnousov, L. A.

ORG: none

TITLE: Radiation polymerization of fluoroolefins

SOURCE: Simpozium po radiatsionnoy khimii polimerov. Moscow, 1964. Radiatsionnaya khimiya polimerov (Radiation chemistry of polymers); doklady simpoziuma. Moscow, Izd-vo Nauka, 1966, 109-114

TOPIC TAGS: radiation polymerization, halogenated organic compound, polymerization kinetics, reaction mechanism

ABSTRACT: Results of the authors' previously published studies on radiation polymerization of unsaturated fluorine-containing compounds are reviewed, explaining certain characteristics of the process associated with the effects of the electronegative fluorine atom, heterogeneous process conditions and radiolysis products. Tetrafluoroethylene is distinguished by its rapid polymerization under ionizing irradiation, with complete monomer conversion in three hours at -78°C in liquid phase polymerization with 10 rad/sec radiation, and in ten minutes at $+20^{\circ}\text{C}$. The yield of 7×10^6 molec/100ev is the highest known for radiation chemical reactions.

Card 1/2